



Patient Name _____
Date of Birth _____
Gender at Birth ☐ Male ☐ Female
Address _____
City, State, Zip _____
Phone _____
Email _____
Primary Insurance _____
Secondary Insurance _____

Prescriber Name _____
NPI # _____ DEA # _____
State Lic # _____ UPI # _____
Practice Name _____
Address _____
City, State, Zip _____
Office Contact _____
Office Contact Phone _____
Office Contact Email _____

CLINICAL INFORMATION

Primary Diagnosis/ICD10 _____ Current Therapy _____
Diagnosis Date _____ Current Directions _____
Prior Therapies/Duration of Use _____
Access Type ☐ Peripheral IV ☐ PICC ☐ Implant Port ☐ Broviac/Hickman Last Dose _____ Next Dose _____
Negative HBV ☐ Yes ☐ No Date of Test: _____ Weight _____ Height _____
Negative TB ☐ Yes ☐ No Date of Test: _____ Allergies _____

PRESCRIPTION INFORMATION

Medication	Avsola®
Directions	☐ Moderately to Severely Active Rheumatoid Arthritis Dose: 3mg/kg Total dose: _____ mg Quantity: <u>1</u> Refills: <u>1 year</u> Directions: _____ mg intravenously on week 0, 2, and 6 then every 8 weeks Administration Rate: ☒ Per Prescribing Information, as tolerated ☐ Other: _____ ☐ Active Ankylosing Spondylitis Dose: 5mg/kg Total dose: _____ mg Quantity: <u>1</u> Refills: <u>1 year</u> Directions: _____ mg intravenously on week 0, 2, and 6 then every 6 weeks Administration Rate: ☒ Per Prescribing Information, as tolerated ☐ Other: _____ ☐ Active Psoriatic Arthritis ☐ Moderately to Severely Active Crohn's Disease ☐ Chronic Severe Plaque Psoriasis ☐ Moderately to Severely Active Ulcerative Colitis Dose: 5mg/kg Total dose: _____ mg Quantity: <u>1</u> Refills: <u>1 year</u> Directions: _____ mg intravenously on week 0, 2, and 6 then every 8 weeks Administration Rate: ☒ Per Prescribing Information, as tolerated ☐ Other: _____
Pre-Medications	☒ Acetaminophen 650mg 30 minutes prior to infusion PO ☒ Diphenhydramine 25mg 30 minutes prior to infusion PO ☐ Solu-Cortef _____ mg slow IVP ☐ Solu-Medrol _____ mg slow IVP Administered: ☐ Pre ☐ Halfway ☐ Upon Completion
Anaphylaxis Orders and Medications	Orders: 1. Stop Infusion 2. Call 911 and prescribing physician 3. Administer medications per protocol Diphenhydramine 50mg/mL slow IVP/IM Administer 12.5mg (wt. <15kg), Administer 25mg (wt. 15-30kg) Administer 50mg (wt. > 30kg) Qty 1 Epinephrine 1mg/mL – Administer ____ mg (wt. < 15kg) IM, Administer 0.15mg (wt. 15-30kg) IM, Administer 0.3mg (wt. > 30kg) IM Qty: 2 Sodium Chloride 0.9% – Administer 250mL IV at KVO (20ml/hr)
Flushing Protocol	☒ Sodium Chloride 0.9% 5-10mL pre and post medications ☐ Heparin _____ Units/mL _____ mL as needed
Pump and Ancillary Supplies	☒ Pump and supplies as needed for administration and appropriate disposal of infusion materials
Skilled Nursing Visits	☒ As needed for IV access, administration and appropriate clinical monitoring

In order for brand name product to be dispensed when generic is available, prescriber must handwrite "Dispense As Written" to prohibit substitution. By signing below, I certify that the above therapy is medically necessary and that the provided medical information is accurate to the best of my knowledge. I authorize Primera Specialty Infusion to act as an agent to initiate and execute the insurance prior authorization process for this prescription and any future fills of the same prescription for patient listed above. I understand that I can revoke this designation by providing notice to Primera Specialty Infusion. Mail 795 Primera Blvd, Ste 1011, Lake Mary, FL 32746 Phone (689) 303-3338 Fax (689) 303-3240

Prescriber Signature _____ Date _____